

AD\_\_\_\_\_

Award Number: W81XWH-05-1-0323

TITLE: Role of Rac GTPases in Chemokine-Stimulated Breast Carcinoma

PRINCIPAL INVESTIGATOR: Symons, Marc H., Ph.D.

CONTRACTING ORGANIZATION: North Shore University Hospital  
Manhasset, NY 10030

REPORT DATE: July 2006

TYPE OF REPORT: Annual

PREPARED FOR: U.S. Army Medical Research and Materiel Command  
Fort Detrick, Maryland 21702-5012

DISTRIBUTION STATEMENT: Approved for Public Release;  
Distribution Unlimited

The views, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy or decision unless so designated by other documentation.

REPORT DOCUMENTATION PAGE				Form Approved OMB No. 0704-0188	
Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing this collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to Department of Defense, Washington Headquarters Services, Directorate for Information Operations and Reports (0704-0188), 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number. <b>PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ADDRESS.</b>					
1. REPORT DATE (DD-MM-YYYY) 01-07-2006		2. REPORT TYPE Annual		3. DATES COVERED (From - To) 1 JUL 2005 - 30 JUN 2006	
4. TITLE AND SUBTITLE Role of Rac GTPases in Chemokine-Stimulated Breast Carcinoma				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER W81XWH-05-1-0323	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S) Symons, Marc H., Ph.D.  E-Mail: <a href="mailto:msymons@nshs.edu">msymons@nshs.edu</a>				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES)  North Shore University Hospital Manhasset, NY 10030				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) U.S. Army Medical Research and Materiel Command Fort Detrick, Maryland 21702-5012				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION / AVAILABILITY STATEMENT Approved for Public Release; Distribution Unlimited					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT: CXCR4 is highly expressed in breast carcinoma cells and is essential for breast cancer metastasis to the lung. CXCR4 is the receptor for CXCL12, a chemokine that is enriched in organs that are targeted by metastatic breast cancer, such as lung and liver. The molecular mechanisms of CXCR4-mediated breast cancer metastasis however are poorly understood. In this project we test the hypothesis that Rac proteins are essential for CXCR4-mediated breast carcinoma cell proliferation and survival, thereby contributing to breast cancer metastasis. The Rac proteins examined comprise Rac1, Rac1b and Rac3. In Task 1, we investigate the role of Rac proteins in CXCL12-regulated breast carcinoma cell proliferation and survival <i>in vitro</i> . In Task 2, we will determine the contribution of these Rac proteins to breast cancer metastasis <i>in vivo</i> . These approaches should allow us to validate Rac-controlled signaling proteins as novel therapeutic targets for metastatic breast cancer. The research performed in this first year of funding largely pertains to Task 1. We showed that CXCL12 stimulates Rac1 activity in breast carcinoma cells, suggesting that Rac1 is likely to play a role in CXCL12- mediated functions in these cells. We also discuss efforts to specifically downregulate the Rac1b splice form.					
15. SUBJECT TERMS cancer cell biology, metastasis models, chemokines, Rac1, Rac1b, Rac3, CXCL12, CXCR4, proliferation, survival, RNA interference					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT	18. NUMBER OF PAGES	19a. NAME OF RESPONSIBLE PERSON
a. REPORT	b. ABSTRACT	c. THIS PAGE			USAMRMC
U	U	U	UU	15	19b. TELEPHONE NUMBER (include area code)

## Table of Contents

<b>Cover.....</b>	<b>1</b>
<b>SF 298.....</b>	<b>2</b>
<b>Introduction.....</b>	<b>4</b>
<b>Body.....</b>	<b>5</b>
<b>Key Research Accomplishments.....</b>	<b>11</b>
<b>Reportable Outcomes.....</b>	<b>12</b>
<b>Conclusions.....</b>	<b>13</b>
<b>References.....</b>	<b>14</b>
<b>Appendices.....</b>	<b>15</b>

## INTRODUCTION

This project examines the roles of the small GTPases Rac1 and Rac3 in CXCR4-mediated metastasis of breast carcinoma cells. CXCR4 is highly expressed in breast carcinoma cells and is the receptor for CXCL12, a chemokine that is produced in abundance in organs that are targeted by metastatic breast cancer, such as lung and liver. Rac1b is a splice form of Rac1 that is constitutively active and expressed in breast cancer tissue [1] and therefore the specific role of this splice form in CXCR4-mediated breast carcinoma metastasis will be examined separate from the role of Rac1 itself. The first aim is to determine the specific roles of Rac1, Rac1b and Rac3 in CXCL12-stimulated carcinoma cell survival and proliferation. The second aim is to examine the roles of Rac1, Rac1b and Rac3 in two animal models of breast cancer metastasis, the first using orthotopic mammary fat pad-implanted human breast carcinoma cells (spontaneous metastasis) and the second tail vein-injected cells (experimental metastasis). These approaches should allow us to validate the potential of Rac proteins and signaling elements controlled by these proteins as novel drug targets for metastatic breast cancer.

### Note:

Tasks 1 and 2, as outlined in the Body of the report below, refer to the Tasks described in the revised Statement of Work, effective June 8, 2006.

## BODY

### **Task 1. To determine the role of Rac proteins in CXCL12-regulated functions in breast carcinoma cells *in vitro*.**


Task 1a: Determine the contribution of Rac1 and Rac3 to CXCL12-stimulated proliferation and survival of breast carcinoma cells.

Choice of cell lines. To ensure that our results can be generalized, we will use two breast carcinoma cell lines. In our application we proposed to employ the DU4475 and MDA-MB-231 cell lines. DU4475 cells express a very high level of CXCR4 mRNA [2] and protein (our unpublished data). MDA-MB-231 cells express a 5-fold lower level of CXCR4 mRNA and protein, but their metastatic potential has been extensively characterized [2;3].

Although the DU4475 cell line was derived from a metastatic nodule from a patient with advanced breast cancer, recent observations have shown that these cells are not metastatic when injected orthotopically in the mouse mammary fat pad (H. Nakshatri, U. Indiana, personal communication), makes them inappropriate for our studies.


Surprisingly, MDA-MB-231 cells that we obtained from ATCC showed negligible expression levels of CXCR4, whereas DU4475 cells displayed robust expression (Fig. 1). We found similar low levels of CXCR4 in cells that we obtained from two colleagues (data not shown). Interestingly, work from the group of H. Nakshatri had shown that MDA-MB-231 cells tend to rapidly lose cell surface expression of CXCR4 when cultured *in vitro*, but regain expression after implantation in the mouse mammary fat pad [4]. Thus, we contacted Dr. Nakshatri to request a vial of mammary fat pad-isolated MDA-MB-231 (termed T231). After a delay of several months we obtained these cells from Dr. Nakshatri. The considerable lag time was due to the fact that Dr. Nakshatri had to reisolate fresh cells after orthotopic transplantation. Determination of the CXCR4 expression levels showed that the T231 that we obtained from Dr. Nakshatri indeed displayed significantly higher expression levels than the other cultures of MDA-MB-231 that we have tested (Fig. 1).

CXCL12 significantly stimulates the proliferative behavior of T231 cells, both in the presence of full serum and low serum (1%) conditions (Fig. 2). We are currently examining whether short interfering RNA- (siRNA)-mediated depletion of Rac1 or Rac3 inhibits CXCL12-stimulated proliferation of T231 cells. A technical problem that we have experienced with these functional studies is the variability in the quality of the CXCL12 preparations that we have obtained from Peprotech, a company with a very good reputation in the chemokine field. We recently have purchased CXCL12 from R&D and are currently testing this preparation.



**Figure 1: Flow cytometric analysis of CXCR4 surface expression in human breast carcinoma cell lines.** Shown are the isotype control (dark gray area) and CXCR4-PE (light gray area) populations. Percentage of cells above isotype control background are indicated. Results shown are representative of 4 independent experiments.

**Methods:** Cells were either counted ( $5 \times 10^5$  cells/condition) and spun down (DU4475 suspension) or trypsinized (trypsin neutralized with trypsin inhibitor), counted ( $5 \times 10^5$  cells/condition), and spun down (MD231, TMD231 adherent). Cells were washed once with FACS buffer (1% BSA in PBS with 0.1%  $\text{NaN}_3$ ), stained with either an isotype control antibody (mouse IgG2b-PE, clone 133303, R&D Systems) or CXCR4 antibody (mouse monoclonal anti-human CXCR4-PE, clone 44717, R&D Systems) for 20 min at room temperature, and then washed two times with FACS buffer. The cells were fixed with 1% formaldehyde in PBS and stored at  $4^\circ\text{C}$  until analysis (within 1-2 days after staining) on a FACS Calibur (BD Biosciences). Approximately 10,000 events were analyzed for each cell line/antibody.




**Figure 2: CXCL12 stimulates the proliferation of T231 cells.** Cells were plated at a density of  $1 \times 10^4$  cells/well in 96-well plates with 6 replicates/condition in the presence of either 10% or 1% FBS and the presence or absence of 100 ng/ml CXCL12 (Peprotech). Cell proliferation was measured using the SRB colorimetric assay (Skehan, P et.al. J.Natl.Cancer Inst., 82: 1107-1112, 1990). The data shown are the mean of 6 wells ( $\pm$  SEM).

**Task 1b:** Determine whether Rac1 and Rac3 are activated by CXCL12 in breast carcinoma cell lines.

We examined whether CXCL12 activates Rac1 in breast carcinoma cells using a well-established method that allows the pulling down of activated Rac proteins from cell lysates [5]. We found that CXCL12 indeed enhances Rac1 activation in T231 cells within

then minutes of application of CXCL12 (Fig. 3). We are still optimizing these results, by trying to decrease the background signal in the absence of serum and also plan to perform a detailed kinetic analysis of CXCL12-mediated Rac1 activation.




To date, there are no antibodies that specifically recognize Rac3 commercially available (we hitherto have employed quantitative PCR to verify siRNA-mediated knock-down). We therefore decided to generate Rac3-specific monoclonal antibodies in collaboration with Dr. A. Campos (Transduction Labs). Unfortunately, these attempts have not yielded satisfactory antibodies. We recently heard that Dr. A. Cox (University of North Carolina at Chapel Hill) has generated polyclonal antibodies against the unique C-terminal domain of Rac3 and plan to collaborate with her to examine whether CXCL12 also activates endogenous Rac3.


**Task 1c:** Determine the contribution of Rac1b to CXCL12-stimulated proliferation and survival of breast carcinoma cells.

Rac1b is a splice form of Rac1 that differs from the main splice form of Rac1 by an insert of 19 codons, just C-terminal from the switch II region [1;6;7]. Notably, Rac1b is expressed in breast cancer tissue [1]. Rac1b is constitutively active and does not bind to RhoGDI [1;7;8]. A critical role for Rac1b in breast cancer progression is supported by recent evidence that Rac1b is induced during malignant transformation of mammary epithelial cells and causes oxidative damage to DNA and genomic instability [9].


To evaluate the role of Rac1b in CXCL12-mediated functions in breast carcinoma cells, we first generated and characterized two independent siRNAs targeting Rac1b. We examined the effects of transient transfection of these siRNAs in ZR-75 breast carcinoma cells, since these cells express higher levels of Rac1b than all other breast carcinoma cell lines that we have examined (including MDA-MB-231, MDA-MB-361 and T47D). Although the protein expression level of Rac1b is much lower than that of Rac1 (the major splice form of Rac1) (Fig. 4, middle panel), its level of activation is significantly higher (Fig. 4, top panel). This observation is consistent with the notion that Rac1b is constitutively active [7;8] and underlines the potential importance of Rac1b.



Transient transfection of either oligo directed against Rac1b caused inhibition of Rac1b expression, albeit to different extents, without affecting the expression of Rac1 (Fig. 5).



We have started to examine the role of Rac1b in a number of functions that are also regulated by Rac1, including the formation of lamellipodia and cell proliferation. Surprisingly, although depletion of Rac1b inhibits lamellipodia in proportion to the level of inhibition of Rac1b expression (Fig. 6A), the less potent oligo (Rac1b-1) has a strong inhibitory effect on cell proliferation, whereas the more potent oligo does not significantly affect cell proliferation at all (Rac1b-2) (Fig. 6B). These preliminary data suggest that at least one of the two oligos displays off-target effects, i.e. degradation of mRNAs that show only partial sequence identity with the region targeted in Rac1b. We therefore recently ordered two additional oligos and plan to examine their depletion potential and functional effects.




**Figure 6: Functional analysis of Rac1b in ZR75 cells.** **A.** Effect of Rac protein depletion on lamellipodia formation. Lamellipodia were quantified as the total length of lamellipodial edge divided by the circumference of the cell. At least 20 single cells were traced for each condition. Error bars denote the SEM. Transfection of ZR75 cells was performed as in Fig. 5. 72 hrs later, the cells were seeded on coverslips, serum starved overnight and then stimulated with 1 nM heregulin (R&D Systems, human heregulin1- $\beta$ 1 EGF domain) for 20 min. The actin cytoskeleton was visualized using Rhodamine-phalloidin and lamellipodia formation was quantified as previously described [10]. **B.** Effect of Rac protein depletion on cell proliferation. ZR75 cells were transfected as in (A). Within 24 hrs of transfection, the cells were plated at a density of  $1 \times 10^4$  cells/well in 96-well plates. Cell proliferation was determined using the SRB method (see Fig. 2). Data shown are the average of 6 wells ( $\pm$  SEM) and are representative of 3 independent experiments.

## **Task 2. To determine the contribution of Rac1, Rac1b and Rac3 to CXCL12/CXCR4-mediated breast cancer metastasis.**

### **Task 2a: Construction and production of viruses for retroviral transfection of shRNA.**

We have purchased two retroviral plasmids containing independent small hairpin RNAs (shRNA) targeting Rac1 from Open Biosystems. We first tested the knock-down efficiency of these plasmids using transient transfection in 293T cells, which display a very high transfection efficiency. Only one of the two shRNAs yielded significant reduction in Rac1 protein levels (Fig. 7). We therefore are in the process of generating additional shRNA plasmids based on the Rac1-directed siRNA sequences that we have shown to be very efficient [10].

In the mean time, to test the efficiency of the commercial plasmids using retroviral transduction, we attempted to produce retroviral particles in HEK293-based (LinX) or NIH-3T3-based (PA317) packaging cells. Unfortunately, the plasmids did not yield any retroviral particles, although a control plasmid (pBABEpuro) produced the expected number of particles. Thus, as an alternative strategy, once we have obtained a more efficient shRNA plasmid, we will use transient transfection of this plasmid in T231 cells, and select pools of knock-down cells. In the event that this procedure would not yield



sufficient inhibition of Rac1 expression, we will try retroviral vectors that use a different promoter, such as CMV or EF-1 $\alpha$ .

## KEY RESEARCH ACCOMPLISHMENTS

CXCL12 stimulates Rac1 activity in MDA-MB-231 breast carcinoma cells. CXCL12 also stimulates the proliferation of these cells.

REPORTABLE OUTCOMES  
**None**

## CONCLUSION

Research carried out during the first year of funding has been hampered by a large number of technical difficulties, including the difficulty in maintaining significant CXCR4 expression levels in breast carcinoma cells *in vitro*, problems with the generation of retroviral particles and unexpected off-target effects of Rac1b-targeting oligos. These obstacles have significantly slowed us down in completing the respective tasks that we set out to accomplish. We are currently proceeding as planned however.

Nevertheless, we were able to show that CXCL12 stimulates Rac1 activity in MDA-MB-231 breast carcinoma cells, suggesting indeed that Rac1 is likely to play a role in CXCL12-mediated functions in these cells, including cell proliferation and cytoskeletal organization. Based on these first results, we feel confident that we will be able to complete Tasks 1d and 1e (identification of candidate CXCL12-stimulated signaling elements and their role in CXCL12-stimulated proliferation and survival of breast carcinoma cells) largely within the proposed time frame. We also hope that we will be able to obtain reliable siRNA oligos to accomplish Task 1c (determination of the contribution of Rac1b to CXCL12-stimulated functions, albeit with a delay).

## REFERENCES

1. Schnelzer A, Prechtel D, Knaus U, Dehne K, Gerhard M, Graeff H, Harbeck N, Schmitt M, Lengyel E: Rac1 in human breast cancer: overexpression, mutation analysis, and characterization of a new isoform, Rac1b. *Oncogene* 2000, 19:3013-3020.
2. Muller A, Homey B, Soto H, Ge N, Catron D, Buchanan ME, McClanahan T, Murphy E, Yuan W, Wagner SN, Barrera JL, Mohar A, Verastegui E, Zlotnik A: Involvement of chemokine receptors in breast cancer metastasis. *Nature* 2001, 410:50-56.
3. Tsai MS, Shamon-Taylor LA, Mehmi I, Tang CK, Lupu R: Blockage of heregulin expression inhibits tumorigenicity and metastasis of breast cancer. *Oncogene* 2003, 22:761-768.
4. Helbig G, Christopherson KW, Bhat-Nakshatri P, Kumar S, Kishimoto H, Miller KD, Broxmeyer HE, Nakshatri H: NF-kappaB promotes breast cancer cell migration and metastasis by inducing the expression of the chemokine receptor CXCR4. *J Biol Chem.* 2003, 278:21631-21638.
5. Azim AC, Barkalow KL, Hartwig JH: Determination of GTP loading on Rac and Cdc42 in platelets and fibroblasts. *Methods Enzymol.* 2000, 325:257-263.
6. Jordan P, Brazao R, Boavida MG, Gespach C, Chastre E: Cloning of a novel human Rac1b splice variant with increased expression in colorectal tumors. *Oncogene.* 1999, 18:6835-6839.
7. Matos P, Collard JG, Jordan P: Tumor-related alternatively spliced Rac1b is not regulated by Rho-GDP dissociation inhibitors and exhibits selective downstream signaling. *J Biol Chem.* 2003, 278:50442-50448.
8. Singh A, Karnoub AE, Palmby TR, Lengyel E, Sondek J, Der CJ: Rac1b, a tumor associated, constitutively active Rac1 splice variant, promotes cellular transformation. *Oncogene.* 2004, 23:9369-9380.
9. Radisky DC, Levy DD, Littlepage LE, Liu H, Nelson CM, Fata JE, Leake D, Godden EL, Albertson DG, Nieto MA, Werb Z, Bissell MJ: Rac1b and reactive oxygen species mediate MMP-3-induced EMT and genomic instability. *Nature* 2005, 436:123-127.
10. Chan AY, Coniglio SJ, Chuang YY, Michaelson D, Knaus UG, Philips MR, Symons M: Roles of the Rac1 and Rac3 GTPases in human tumor cell invasion. *Oncogene* 2005, 24:7821-7829.

APPENDICES  
**none**